

PFAS and their impacts on human health: An example of a study of PFAS exposure and blood glucose control in older American adults with type 2 diabetes

Eloïse Brosset, Doctoral student

*Axe Diabetes Obesity Cardiovascular complication– Département community health
Sherbrooke University Hospital Research Center, CRCHUS (Canada)*

Diabetes key fact :

- 2014 : 8,5 % of adult diabetes
- 2021 : 10 % of adult diabetes
- 1,5 millions of deaths in 2019



Diabetes and ageing


5 years less life expectancy

PFAS

5000 to 10,000 compounds

PFHxS (C=6)	<p>The structure shows a six-carbon chain. The first carbon is bonded to one hydrogen and five fluorines. The second and third carbons are each bonded to two fluorines. The fourth and fifth carbons are each bonded to one fluorine. The sixth carbon is bonded to one fluorine and a sulfonic acid group (-SO₃H).</p>
PFOS (C=8)	<p>The structure shows an eight-carbon chain. The first carbon is bonded to one hydrogen and seven fluorines. The second and third carbons are each bonded to two fluorines. The fourth and fifth carbons are each bonded to one fluorine. The sixth and seventh carbons are each bonded to two fluorines. The eighth carbon is bonded to one fluorine and a sulfonate group (-SO₃⁻).</p>
PFOA (C=8)	<p>The structure shows an eight-carbon chain. The first carbon is bonded to one hydrogen and seven fluorines. The second and third carbons are each bonded to two fluorines. The fourth and fifth carbons are each bonded to one fluorine. The sixth and seventh carbons are each bonded to two fluorines. The eighth carbon is bonded to one fluorine and a carboxylic acid group (-COOH).</p>
PFNA (C=9)	<p>The structure shows a nine-carbon chain. The first carbon is bonded to one hydrogen and eight fluorines. The second and third carbons are each bonded to two fluorines. The fourth and fifth carbons are each bonded to one fluorine. The sixth and seventh carbons are each bonded to two fluorines. The eighth carbon is bonded to one fluorine. The ninth carbon is bonded to one fluorine and a carboxylic acid group (-COOH).</p>
PFDeA (C=10)	<p>The structure shows a ten-carbon chain. The first carbon is bonded to one hydrogen and nine fluorines. The second and third carbons are each bonded to two fluorines. The fourth and fifth carbons are each bonded to one fluorine. The sixth and seventh carbons are each bonded to two fluorines. The eighth carbon is bonded to one fluorine. The ninth carbon is bonded to one fluorine. The tenth carbon is bonded to one fluorine and a carboxylic acid group (-COOH).</p>

Sources of PFAS ?



Problematic :

Does the presence of PFAS in the blood influence the chances of achieving good blood glucose control in people aged 65 and over?



National Health and Nutrition Examination Survey

- Bi-annual survey
- Since 1960 and continuous study from 1999
- Examines at least 5000 people each year
- 4575 participants for our study



Statistics :

- Linear mixed models : GLIMMIX procedure in the SAS statistical package (version 9.4. SAS)
- The PFAS distributions categorized into quartiles
- For the association of PFAS with poor glycemic control : Adjusted odds ratio (aOR)

Results :

Distribution of pre- and polyfluoroalkyl substances in US adults aged 65 years and more, NHANES 1999-2018


	Mean (SD) ^a	p25	p50	p75	p90
Overall					
PFDeA	0.36 (0.42)	0.14	0.20	0.40	0.70
PFNA	1.04 (1.20)	0.50	0.74	1.30	2.00
PFOA	3.64 (3.81)	1.87	3.00	4.30	6.30
PFOS	20.14 (22.0)	7.70	14.50	24.90	40.80
PFHxS	2.39 (2.71)	0.90	1.70	2.60	4.80
ΣPFAS	14.8 (21.4)	2.30	5.80	20.70	37.7
Men					
PFDeA	0.38 (0.44)	0.16	0.20	0.40	0.80
PFNA	1.06 (0.95)	0.50	0.80	1.30	1.90
PFOA	3.82 (4.82)	1.80	2.90	4.17	6.40
PFOS	21.91 (25.92)	8.90	15.40	26.10	43.70
PFHxS	2.76 (3.15)	1.20	1.90	3.00	5.30
ΣPFAS	15.31 (23.41)	2.70	5.60	20.37	36.82
Women					
PFDeA	0.34 (0.41)	0.10	0.20	0.40	0.70
PFNA	1.03 (1.43)	0.40	0.70	1.20	2.10
PFOA	3.47 (2.39)	1.87	3.04	4.39	6.29
PFOS	18.37 (16.92)	6.81	13.70	23.40	39.70
PFHxS	1.97 (2.03)	0.90	1.40	2.20	3.70
ΣPFAS	13.92 (18.50)	2.00	5.80	20.60	37.59

Results :

(A) poor glycemic control defined as HbA1c $\geq 7.0\%$ in 65–74 years of age and $\geq 8.0\%$ in order;


(B) poor glycemic control defined as HbA1c $\geq 7.5\%$ in 65–74 years of age and $\geq 8.0\%$ in older.

PFNA



PFHxS

Perfluorohexanesulfonic acid



Results :

Odds ratio (95% confidence intervals) for the associations between pre- and polyfluoroalkyl substances and poor glycemic control in US men aged 65 years and older


	(B)			
	Q1 (Reference)	Q1-Q2	Q2-Q3	≥Q3
Model 1^a				
PFDeA	1	2.26 (0.16-36.70)	2.55 (0.88-7.37)	7.59 (2.61-22.06)
PFNA	1	0.17 (0.07-0.44)	0.57 (0.23-1.41)	1.74 (0.72-4.20)
PFOA	1	0.18 (0.04-0.76)	0.40 (0.11-1.51)	0.33 (0.08-1.32)
PFOS	1	1.56 (0.36-6.72)	1.22 (0.28-5.34)	3.39 (0.74-15.49)
PFHxS	1	6.14 (2.69-14.05)	0.58 (0.24-1.41)	6.47 (2.70-15.46)
ΣPFAS	1	1.22 (0.65-2.27)	0.35 (0.12-1.02)	0.45 (0.11-1.81)
Model 2^b				
PFDeA	1	2.72 (0.17-43.51)	4.44 (1.78-11.07)	6.13 (2.42-15.50)
PFNA	1	0.27 (0.12-0.61)	0.51 (0.24-1.09)	0.90 (0.43-1.88)
PFOA	1	0.16 (0.04-0.62)	0.35 (0.10-1.23)	0.30 (0.08-1.12)
PFOS	1	0.89 (0.24-3.26)	0.94 (0.25-3.47)	2.13 (0.56-8.11)
PFHxS	1	5.46 (2.50-11.93)	0.58 (0.24-1.39)	3.80 (1.73-8.33)
ΣPFAS	1	1.10 (0.62-1.98)	0.33 (0.12-0.86)	0.41 (0.11-1.44)
Model 3^c				
PFDeA	1	2.49 (0.17-36.55)	2.83 (1.32-6.06)	7.11 (3.13-16.13)
PFNA	1	0.52 (0.26-1.05)	1.20 (0.58-2.48)	2.75 (1.30-5.81)
PFOA	1	0.32 (0.11-0.93)	0.31 (0.10-0.97)	0.29 (0.09-0.92)
PFOS	1	1.01 (0.36-2.86)	1.17 (0.36-2.86)	1.25 (0.40-3.93)
PFHxS	1	5.88 (2.91-11.91)	0.83 (0.40-1.75)	5.46 (2.65-11.26)
ΣPFAS	1	1.54 (0.86-2.78)	1.41 (0.62-3.23)	0.85 (0.28-2.59)
Model 4^d				
PFDeA	1	2.42 (0.15-38.35)	2.79 (0.98-7.96)	8.50 (2.98-24.28)
PFNA	1	0.18 (0.07-0.46)	0.62 (0.25-1.52)	1.96 (0.83-4.62)
PFOA	1	0.20 (0.05-0.80)	0.46 (0.12-1.66)	0.39 (0.10-1.49)
PFOS	1	1.45 (0.35-5.97)	1.11 (0.26-4.81)	3.42 (0.76-15.35)
PFHxS	1	5.87 (2.63-13.10)	0.60 (0.25-1.44)	6.84 (2.92-16.02)
ΣPFAS	1	1.24 (0.67-2.29)	0.37 (0.12-1.08)	0.51 (0.13-2.05)

(B) poor glycemic control defined as HbA1c ≥ 7.5% in 65–74 years of age and ≥8.0% in older.



Strengths :

- 1st study to focus on elderly populations and glycemic control in cases of known diabetes
- Large sample size and heterogeneous population in terms of race/ethnicity



Limits :

- Residual confounding
- Causal inference
- Multiple co-exposure
- Treatments not taken into account
- No consideration for food

Conclusion

This is the first study to assess glycemic control in relation to PFAS exposure in older populations with known diabetes.

PFNA

PFHxS

PFDeA (for men)



Poor glycemic control at U.S.
adults aged ≥ 65 years with
known T2DM

Perspectives

- Assessing the effect of mixtures
- Evaluating PFAS and its influence on the oral glucose test in adults of various ages
- Repeat analyses with longitudinal data
- Work with more PFAS for a more global scheme (including alternative PFAS)
- Enable prevention campaigns by formation healthcare professionals to alert people at risk

Acknowledgements



CENTRE DE
RECHERCHE



My thesis supervisors

- Pr Gérard Ngueta



- Pr William Fraser





Thanks for your attention

References

- Brosset E, Ngueta G. Exposure to per- and polyfluoroalkyl substances and glycemic control in older US adults with type 2 diabetes mellitus. *Environ Res.* 2023 Jan 1;216(Pt 3):114697. doi: 10.1016/j.envres.2022.114697. Epub 2022 Nov 2. PMID: 36334831.
- *Diabète.* (s. d.). Consulté 7 juin 2023, à l'adresse <https://www.who.int/fr/news-room/fact-sheets/detail/diabetes>
- Manojkumar Y, Pilli S, Rao PV, Tyagi RD. Sources, occurrence and toxic effects of emerging per- and polyfluoroalkyl substances (PFAS). *Neurotoxicol Teratol.* 2023 May-Jun;97:107174. doi: 10.1016/j.ntt.2023.107174. Epub 2023 Mar 11. PMID: 36907230.