

ENVIRONMENTAL METROLOGY CHALLENGES FOR A BETTER UNDERSTANDING OF THE SOURCES AND TRANSFER OF PFAS IN THE ENVIRONMENT

Anne **TOGOLA**, S.Bristeau, B.Idjaton, C.Coureau, J.Lions, JP. Ghestem

What are the challenges facing environmental monitoring?

✓ Over 10,000 compounds: which ones to track?

Source-specific compounds

Poor information (industrial secrecy / user ignorance)

Temporal evolution of uses (regulatory restrictions + technological developments)

✓ Different pathways of degradation/transfer depending compartments

Need to know degradation pathways

Specificity of different compartments (soil/surface water/groundwater)

To answer which questions? Occurrence?, Risk? Management?

These questions determine **what** to track, **where, how** and associated **analytical performances to reach**.


PFOA example..

End of production 2011 / end of use 2023

aqueous film-forming foam (AFFF)

degradation

Used in



PFOA


Used in

Polymers PVDF, PTFE

Used in

Pesticides ?


- Same compound as source and degradation by product :
 ⇒ need to identify precursors and degradation pathways ..



The PFAS Universe

Less than 200 compounds :

- Identified
- with analytical standards
- with adapted methods



Expectation for PFAS

Most « relevant compounds »

PFAS to identify the source of pollution

PFAS to be remediate

PFAS to monitor in drinking water

...

At low concentrations levels

In all matrix of interest


...

An overview of stakeholder expectations...



What were the main challenges in making steps towards foreseen advancement beyond the state of the art (related to Ambition1)

Methods developed depending on the objectives to reach



What are the challenges facing environmental monitoring?

knowledge challenges :

- New PFAS of interest ?
- Precursor's fate ?
- PFAS pattern related to specific sources ?
- Degradation by-products during remediation ?
- ...

Target quantitative analysis

Semi-globales quantitative analysis

Globales quantitative analysis

Qualitative analysis

Objectives; : Propose and distinguish:

- Methods fit for purpose for research needs ?
- Method fit for purpose for regulation, management ..

Regulatory issues

List of few PFAS (20-28)

Need of low analytical performance (ng/L sub ng/L)


Other methods fit for purpose ??

Target quantitative analysis

Approaches implemented at BRGM

Target quantitative analysis


Constant implementation since 5 years..
based on needs and opportunities ..




- + Robust and reliables methods
- Limitation based on available standards
- Need of several analysis to cover different needs :
from LC/MS/MS to IC/MS/MS and GC/MS/MS

Impact of the targeted list of compounds on data understanding

WWTP A_ 20 PFAS, $\Sigma = 180$ ng/L




WWTP A_ 56 PFAS, $\Sigma = 470$ ng/L




2 urban WWTP on the same areas (5 km)
Different PFAS fingerprints with the PFAS-20
but highly underestimated without 2 compounds..

WWTP B_ 20 PFAS, $\Sigma = 120$ ng/L




WWTP B_ 56 PFAS, $\Sigma = 530$ ng/L



Semi-globales quantitative analysis
Total oxidizable precursors « TOP »

- To have a better picture about the PFAS content
- Considering that main PFAS can be oxidized in a restricted list of PFAS





Zhang et al, EST Letters, 2019


- + More complete picture
- + Assesment of the potential of the « PFAS contamination » to be degraded in « persistant PFAS »
- **Method should be better defined (lack of robustness , list of compounds analyzed, need of data on oxidability of some precursors..)**
 - No information on PFAS pattern
 - should be used in association with targeted analysis

Some examples..

Blue : PFCA/PFSA « 20 DWD »
 Orange : other PFAS oxidisable
 Yellow : other PFAS partially oxidisable ..



TOP assay give a simplified view ... based on short list of compounds





BUT : some compounds are not oxidisable.. or not completely ..

Interesting method:
Easier way to have a better overview
But need more robustness and QA/QC methods


| | concentration ng/L | | 6:2 FTS oxidation efficiency |
|-------------------|--------------------|-----------|------------------------------|
| | before TOP | after TOP | |
| WWTP | 1285 | 126 | 90% |
| WWTP | 391 | | 100% |
| Indus WWTP | 1092 | 229 | 79% |
| Indus WWTP | 24624 | 136 | 99% |
| Indus WWTP | 1062367 | 1572 | 100% |

Global quantitative analysis

- Global analysis of fluorine compounds




Comparison AOF and TF



- ❖ AOF method can explain 1 to and 23% of the TF.
- ❖ Results depend on samples.
- ❖ AOF seems not representative of the fluorinated compounds concentration in the environment.



Comparison AOF and Target analysis



- ❖ Only a very small fraction (<10%) of AOF is explained by targeted LCMSMS analysis (56 PFAS). Except for one sample where LCMSMS was able to explain 74% of the fluorine in the AOF.
- ❖ This value is very low compared to TF.


Qualitative analysis

Which PFAS in AFFF ?



Targeted analysis (56)
identification of 20% of the TF (PFAS)
Up to 18 unknowns ..
Biblio review = 1 candidate
80 % identification (PFAS-57)
And now PFAS 58 ?

But time consuming.. And so many unknown..
And dependant on standard availability ..



Qualitative analysis with HRMS



2024-2025

In 2018 a database of digital samples (NTS fingerprint) of 85 surface waters has been built

Suspect screening applied :

Are a list of compounds (56 PFAS) in my samples?

Comparison of mass fingerprints in samples with « reference compounds fingerprints » in internal database



No QA/QC focused on PFAS in 2018 : risk of sample contamination
Are the extraction /analytical methods adapted to PFAS ?


- ✓ Recovery test (20- 60% recovery)
- ✓ Control of field blanks (only background level)
- ✓ Control of analytical blanks

Caution in data interpretation due to analytical uncertainties, but :

- 27 PFAS detected (list of 56..)
 - 6:2 FTAB detected in 55% of the water sample with a high level of confidence, PFOS in 5% and PFBS in 15%
 - More than 10 different PFAS detected in 10% of the samples
 - Qualitative information !!
 - retrospective quantification is not robust
-
- **Upcoming actions:**
 - identification of potential sources in correlation with the hot spots
 - Confirmation by new sampling campaign should be planned
 - Complementary approach using other PFAS databases to search for more compounds
 - Implementation of a new database based on real products (eg AFFF)



A lot of challenges for the PFAS monitoring
A lot of available tools..



CIC: combustion ion chromatography
TF: Total fluorine TIF: Total inorganic fluorine
TOF: Total organic fluorine EOF: Extractable organic fluorine
TOP Assay: Total Oxydable Precursor assay

But also a lot of needs :

The need to acquire knowledge while making progress on management and regulations

The need to communicate despite scientific uncertainty

The Need to regulate while waiting for new knowledges

= not the same tools / not the same players / not the same barriers!

Need for consultation, transparency on advantages/disadvantages/uncertainties on what we know ... and on what we don't know



Thanks ..

a.togola@brgm.fr

The team...

Babatoundé Idjaton, PhD PROMISCES

Sébastien Bristeau, Emeline Coisy, Charlotte Coureau, LC/MS; LCHRMS

Laura Kastler, analysis CIC

Jean Philippe Ghestem, normalization, AQUAREF

Julie Lions PROMISCES coordinator

...



This project has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement N°101036449

